

be agricultural pests, compared to the cryptic smooth larvae.

I thank Jane Ruffin (Rosemont, PA) for discovering and showing me the population of *Parasa indetermina* and Neal Evenhuis (Bishop Museum) for determining the spec-

imens of *S. macer*, providing helpful bibliographic information and searching the USNM collection for host information.

Marc E. Epstein, *Department of Entomology, MRC 105, Smithsonian Institution, Washington, D.C. 20560, U.S.A.*

PROC. ENTOMOL. SOC. WASH.
99(3), 1997, pp. 586

NOTE

Replacement Names for Western Hemisphere Genera of Doryctinae (Hymenoptera: Braconidae)

Recently I published descriptions of several new genera of Doryctinae from the Western Hemisphere (Marsh 1993. *Contributions of the American Entomological Institute* 28(1): 1-58). Unfortunately, three of these generic names have been previously used and must be replaced with new names. Therefore, I propose the following new generic names.

Cyphodoryctes Marsh, **new name**. Replacement name for *Cyrtonion* Marsh 1993, not *Cyrtonion* Hansen 1989 (see Hansen 1991. *Kongelige Danske Videnskabernes Selskab Biologiske Skrifter* 40: 1-367). The original name *Cyrtonion* was derived from the Greek *kyrton* meaning humpbacked in reference to the mesonotum which anteriorly is swollen and strongly declivous. The

replacement name is derived from the Greek *kyphos* also meaning humpbacked.

Donquickeia Marsh, **new name**. Replacement name for *Quickia* Marsh 1993, not *Quickia* Odhner 1950 (see Vaught 1989. A classification of the living Mollusca. American Malacologists, Inc.).

Whitfieldiellus Marsh, **new name**. Replacement name for *Whitfieldia* Marsh 1993, not *Whitfieldia* Davidson 1882 (see Moore, ed., 1965. *Treatise on invertebrate paleontology. Part H, Brachiopoda. Vols. 1 and 2.* The Geological Society of America and the University of Kansas).

Paul M. Marsh, *P.O. Box 384, North Newton, KS 67117, U.S.A.*